

eRed Folder : [First Hit](#)[Previous Doc](#)[Next Doc](#)[Go to Doc#](#)☐

L1: Entry 232 of 281

File: DWPI

Nov 21, 2007

DERWENT-ACC-NO: 2008-L74864

DERWENT-WEEK: 200869

Copyright ©2011 Thomson Reuters

TITLE: Skin protective formulation e.g. ointment, cream and emulsion for promoting scar restoration, comprises metallothionein, epidermal growth factor, additional skin protective active agent and carrier

INVENTOR: CAO L ; JIANG G ; JIANG L ; ZHOU L

PATENT-ASSIGNEE: HUNAN LIGU BIOLOGICAL TECHNOLOGY LTD (HUNAN)

PRIORITY-DATA: 2006CN-10031677 (May 19, 2006)

PATENT-FAMILY:

PUB-NO

PUB-DATE

LANGUAGE

☐ [CN 101073539 A](#)

November 21, 2007

ZH

APPLICATION-DATA:

PUB-NO

APPL-DATE

APPL-NO

DESCRIPTOR

CN 101073539A

May 19, 2006

2006CN-10031677

INT-CL-CURRENT:

TYPE IPC

DATE

CIPP [A61 K 8/64](#) 20060101CIPS [A61 Q 19/00](#) 20060101

ABSTRACTED-PUB-NO: CN 101073539 A

BASIC-ABSTRACT:

NOVELTY - Skin protective formulation comprises metallothionein (50-200 parts per minute), epidermal growth factor (50000-600000 IU/30 ml), additional skin protective active agent and carrier.

USE - The skin protective formulation such as cream, ointment, facial membrane and emulsion is useful for promoting scar restoration (all claimed), for healing the scar of skin, reducing the scar and pigmentation of the scar.

ADVANTAGE - The compound rapidly heals the scar and improves the skin quality.

ABSTRACTED-PUB-NO: CN 101073539 A
EQUIVALENT-ABSTRACTS:

ORGANIC CHEMISTRY

Preferred Components: The metallothionein is zinc-binding metallothionein extracted from rabbit liver, the epidermal growth factor is recombinant human epidermal growth factor. The dosage of metallothionein is 80-160 ppm, preferably 100-140 ppm, and the dosage of the epidermal growth factor is 100000-600000 IU/30 ml, preferably 200000-500000 IU/30 ml. The additional skin protective active agents are chosen from water-soluble vitamin E, hyaluronic acid powder, sodium heparin, apricot kernel oil and propolis. The carrier is acrylic acid ester/10-30C alkyl acrylic acid ester crosslinking copolymer, pyrrolidone sodium carboxylic acid, squalane and 1,3-butanediol.

The skin protective compound when formulated as cream, comprises (in ml) aqua (30) with freeze dry meal (20). The aqua includes (in weight%) acrylic acid ester/10-30C alkyl acrylic acid ester/10-30C alkyl acrylic acid ester crosslinking copolymer (0.2-0.5), 1,3-butanediol (4-6), glycerin (1-3), pyrrolidone sodium carboxylic acid (0.5-2), vitamin E (0.3-0.7), hyaluronic acid (0.2-0.6), propolis (0.2-0.5), sodium heparin (0.3-0.5), dimethylol hydantoin and 3-iodine-2-propinyl amido butyl formate (0.1-0.3), metallothionein (2-7 ml/20 ml), epidermal growth factor (100000-500000 IU/20 ml), and mannitol (18 ml/20 ml). The skin protective compound when formulated as facial membrane, comprises (in weight%) acrylic acid ester/10-30C alkyl acrylic acid ester/10-30C alkyl acrylic acid ester crosslinking copolymer (0.2-0.5), 1,3-butanediol (4-6), glycerin (1-3), pyrrolidone sodium carboxylic acid (0.5-2), silk peptide powder (0.1-0.3), vitamin E (0.4-0.8), hyaluronic acid (0.2-0.6), dimethylol hydantoin and 3-iodine-2-propinyl amido butyl formate (0.1-0.3), metallothionein (60-160 parts per minute) and epidermal growth factor (150000-500000 IU/20 ml). The skin protective compound when formulated as ointment, comprises (in weight%) 1,3-butanediol (4-6), glycerin (1-3),

hexadecanol (3-5), octodecyl alcohol (1-3), squalane (3-5), stearin tristearin (0.5-2), apricot kernel oil (1-3), vitamin E (0.2-0.5), hyaluronic acid (0.4-0.6), dimethylol hydantoin and 3-iodine-2-propinyl amido butyl formate (0.1-0.3), metallothionein (60-140 parts per minute) and epidermal growth factor (100000-500000 IU/20 ml). The skin protective compound when formulated as emulsion, comprises (in weight%) 1,3-butanediol (2-5), hexadecanol (3-5), octodecyl alcohol (0.5-3), squalane (1-4), stearin tristearin (0.5-2), apricot kernel oil (1-3), vitamin E (0.2-0.5), hyaluronic acid (0.4-0.6), collagen (0.1-0.4), silk peptide powder (0.1-0.3), dimethylol hydantoin and 3-iodine-2-propinyl amido butyl formate (0.1-0.3), metallothionein (60-170 parts per minute) and epidermal growth factor (100000-500000 IU/20 ml).

DERWENT-CLASS: A96 B04 D21

CPI-CODES: A12-V04C; B03-H; B04-B01C; B04-C02E2; B04-C03B; B04-H06A0E; B04-N02; B05-A03A4; B07-D03; B07-D09; B10-A07B; B10-D03; B10-E04C; B10-J02; B12-M02B; B12-M03; B14-N17; B14-R01; B14-S18; D08-B01D1;

[Previous Doc](#)

[Next Doc](#)

[Go to Doc#](#)